

Analysis of the Bohm criterion for two-ion-species plasmas using PARASOL

S. Azuma^{1*}, A. Fukuyama¹, T. Takizuka²

¹ *Department of Nuclear Engineering, Kyoto University, Yoshida, Kyoto 606-8501, Japan*

² *Japan Atomic Energy Agency, 801-1 Mukoyama, Naka 311-093, Japan*

* *Email: azuma@p-grp.nucleng.kyoto-u.ac.jp*

The divertor plays essential roles in tokamak reactors for heat removal, ash exhaust, and impurity shielding. It is important to study the physics of sheath formation at a divertor plate exposed to plasmas and the interaction between plasma and wall. The Bohm criterion for a single-ion-species plasma forming a stable sheath is given by the incident ion flow velocity equal to the sound velocity. For multi-ion-species plasmas such as DT plasmas in fusion devices, however, the criterion is not still fully understood. The usual analytical procedure does not lead to a unique solution but a relation between the ion flow velocities [1]. To clarify the Bohm criterion for two-ion-species plasmas, Tskhakaya once studied it preliminarily with particle simulations taking into account all kinetic effects [2]. In the present paper, we evolutionally carry out particle simulations on this subject with a one-dimensional particle code PARASOL [3]. We found that when collisionality $L_{||}/l_{\text{mfp}}$ ($L_{||}$: system size along the magnetic field, and l_{mfp} : mean free path) is low, $u_2/u_1 \approx \sqrt{m_1/m_2}$ at the entrance of magnetic presheath (u_s : ion flow velocity, m_s : ion mass and the index s represents ion species), while $u_1 = u_2$ when collisionality becomes high.

We will report the dependence of the Bohm criterion on collisionality and other plasma parameters. We will also present the peculiar sheath physics for two-ion-species plasmas.

[1] K.-U. Rieman, IEEE Trans. Plasma Sci. **23** (1995) 709.

[2] D. Tskhakaya, S. Kuhn, Y. Tomita, Contrib. Plasma Phys. **46** (2006) 649.

[3] T. Takizuka, M. Hosokawa, K. Shimizu, Trans. Fusion Technol. **39** (2001) 111.